

BAY AREA COLON & RECTAL SURGEONS

A DIVISION OF BASS MEDICAL GROUP

Drs. Ran Kim, Brian McGuinness and Negar Salehomoum

365 Lennon Lane # 290, Walnut Creek CA 94598

925-274-9000

INSTRUCTIONS FOLLOWING ANORECTAL SURGERY

PAIN AND DISCOMFORT

There is usually some bleeding and discomfort associated with bowel movements in 1-2 weeks after surgery. However, if you experience excessive bleeding, increasing pain or develop a high fever, please call our office. It is also common for a hard, uncomfortable lump to form, as well pressure sensations in your rectum. This is a normal occurrence that will heal gradually as your body adjusts to the trauma undergone during surgery. If you need to refill your pain medication, please call the office to request a written prescription from your doctor. Please allow 48 hours for your request to be processed.

URINARY SYMPTOMS

Some patients have experienced difficulty in urinating after surgery. This can usually be relieved by warm sitz baths.

CONSTIPATION

When you arrive home, take one (1) heaping teaspoon of **METAMUCIL** and mix it up with water or juice at least 2 times a day. Continue this for one month. It is common not to have a bowel movement 2-3 days after surgery. If this happens, take 2-3 tablespoons of **MILK OF MAGNESIA**. If this does not produce results, call your physician. It is very important not to strain during bowel movements as the area is very delicate.

DIET

It is important that you stay on a high fiber diet. Drink at least eight (8) glasses of water a day and avoid excessive quantities of milk products to prevent constipation and hard bowel movements.

ACTIVITY

Do not do any heavy lifting or straining the week following surgery. Try to avoid driving as much as possible the first week.

LOCAL CARE OF THE WOUND

Soak in a bathtub for 15 minutes 3-4 times a day to help clean the area, decrease swelling and relieve muscle spasms. To soothe the area, apply **BALNEOL LOTION** with a damp washcloth. Afterwards to absorb the moisture and secretions of the incision site, gently tuck a piece of gauze or cotton. Please do not use any tape.